

ASC PREOP CONTINUATION OF MEDICATIONS – continue **all meds except:**

ANTICOAGULANTS (including aspirin) – instructions on holding anticoagulants need to come from the prescriber, the surgeon, or a consultant

NSAIDs: Hold for 4 days (unless otherwise instructed by the surgeon)

INSULINS: Continue as normal until the morning of procedure:

- If BG > 140, take one-half of AM dose of long-acting insulins (LANTUS, LEVEMIR, TOUJEO, TRESIBA)
- Pumps – suggest decreasing the basal rate by one-half (patient may override this instruction)
- Hold all other insulins

ORAL DIABETIC MEDS: Hold the day of surgery

INJECTABLE NON-INSULIN DIABETIC MEDS: For those on once daily dosing, just hold the day of surgery. For those on weekly dosing, hold for 7 days whenever possible

GLP1 Agonists	Trade Name	Dosing
Exanetide	BYETTA	SQ Daily
	BYDUREON	SQ Weekly
Semaglutide	WEGOVY	SQ Weekly
	OZEMPIC	SQ Weekly
Liraglutide	RYBELSUS	PO Daily
	VICTOZA	SQ Daily
Dulaglutide	SAXENDA	SQ Daily
	TRULICITY	SQ Weekly
Tirzepatide	MOUNJARO	SQ Weekly

DIURETICS – hold the day of surgery if no history of CHF

NARCOTIC LIMITERS: Can continue as normal for minor surgery, otherwise consult anesthesia for instructions

Narc Modifier	Trade Name	Dosing	Frequency	Use
Buprenorphine	BELBUCA	ODT/SL	Daily	Narc control
	BRIDAXI 8-32	ODT/SL	Weekly	Narc control
	BRIDAXI 64-128	SQ	Monthly	Narc control
	BUTRANS	Patch	Weekly	Narc control
Buprenorphine Naloxone combo	SUBLOCADE	SQ	Monthly	Narc control
	SUBOXONE	ODT/SL	Daily	Narc control
Naltrexone	ZUBSOLV	ODT/SL	Daily	Narc control
	REVIA	PO	Daily	Narc control
Naltrexone with antidepressant	VIVITROL	IM	Monthly	Alcoholism
	CONTRAVE	PO	Daily	Weight loss

HERBALS AND VITAMIN E:

- Ideally hold for 7 days (will not cancel case if this is not possible)
- Cannabis in all forms – Do not consume two hours before arrival

Type-A MAO Inhibitors for depression: Hold for 7 days, but only under psychiatric supervision

MARPLAN, NARDIL, PARNATE